

Maternal and fetal outcomes of home births in Western Sudan

Ahmed Abdelkarim Ahmed Abdallah^{1,2,3} Nawal Ali Ahmed Mohamed^{1,4,5} Ahmed Amin Mohammed^{1,6,7} Khalid Nasralla Habeebulla Hashim⁸ Hussain Gadelkarim Ahmed^{9,10}

¹Department of Obstetrics and Gynecology, Faculty of Medicine, University of Kordofan, NK, El-Obeid, Sudan.

²Alfayhaa Obstetrics, Gynecology, and Laparoscopy Surgery Clinic, El-Obeid, Sudan.

³Sudan medical specialization board, training committee, Sudan.

⁴Sheikan College El-Obeid, Sudan.

⁵Arous Elrimal Obstetrics, Gynecology, and Laparoscopy Surgery Clinic, El-Obeid, Sudan.

⁶Highcare specialized clinic. El-Obeid, Sudan.

⁷Farah fertility center, El-Obeid, Sudan.

⁸Department of Obstetrics and Gynecology – Faculty of Medicine, Qassim University, Buraydah, Saudi Arabia

⁹Prof Medical Research Consultancy Center- MRCC, El-Obeid, Sudan.

¹⁰Department of Histopathology and Cytology, FMLS, University of Khartoum, Sudan.

Abstract

Background: In light of the ongoing Sudanese crisis, many pregnant women are having trouble getting the medical care they seek. Thus, the current study sought to evaluate the effects of home births in Western Sudan on both mothers and their newborns. **Methodology:** From January to December 2023, in the Obstetrics and Gynecology teaching hospital in El-Obeid, North Kordofan State, Sudan, we conducted a descriptive longitudinal investigation for this study. This study has recruited approximately 224 participants. **Results:** The most common maternal outcomes observed in this group of patients were postpartum hemorrhage, followed by puerperal sepsis and eclampsia, accounting for 14.3 percent, 13%, and 9 percent of cases, respectively. Fresh stillbirth accounted for the majority of cases, while macerated stillbirth, fetal distress, and birth trauma accounted for 28%, 17%, 15%, and 9% of the cases, respectively. **Conclusion:** Home birth is widespread in Sudan; however, it might harm the mother and baby. Postpartum hemorrhage, puerperal infection, and eclampsia are the main maternal consequences. Fresh stillbirth, macerated stillbirth, fetal distress, and delivery trauma are common fetal outcomes. Prioritize maternity and prenatal care system improvements.

Keywords: maternal, home birth, pregnancy, fetal outcomes, Sudan.

Correspondence to: Ahmed Abdelkarim Ahmed Abdallah Email: ahsumemo@yahoo.com

Orcid: [0009-0009-3182-3978](https://orcid.org/0009-0009-3182-3978)

Introduction

Maternal mortality, stillbirths, and neonatal mortality cause over 5 million deaths each year. These three categories of deaths are often investigated separately, despite the fact that they share common causes and can be handled using similar approaches [1]. Many women in their prime sadly died early as a result of negligence and avoidable situations throughout the critical stages of pregnancy, labor, and the postoperative period. This is a distressing issue in maternal mortality that affects one woman every two minutes, or 800 women each day [2, 3].

Maternal mortality, as defined by the prestigious World Health Organization (WHO), includes the untimely death of a woman at any stage of her pregnancy, regardless of location or duration, or within 42 days after delivering. This tragic outcome is attributed to circumstances that are inextricably linked to or exacerbated by the pregnancy itself, as well as its management, with the exception of serendipitous and unintended causes [4].

Nevertheless, nations impacted by armed conflict perceive a greater incidence of maternal mortality compared to those that are politically stable [5]. Antenatal care significantly impacts crucial health indices, such as maternal and

neonatal mortality rates, by promptly diagnosing and managing pregnancy-related problems. The political instability and military violence in Sudan have had a profound impact on the health sector, leading to dire consequences for the health of pregnant women. The majority of women residing in remote villages in Sudan rely on traditional birth attendants to deliver their infants. This can be potentially hazardous or unclean. Hence, the objective of this study was to evaluate the maternal and fetal outcomes of home deliveries in Western Sudan. Nevertheless, nations impacted by armed conflict see elevated rates of maternal mortality compared to those that are politically stable. Hence, the objective of this study was to evaluate the maternal and fetal outcomes of home deliveries in Western Sudan.

Materials and Methods

This study is a descriptive longitudinal inquiry that took place at the Obstetrics and Gynecology hospital in El-Obeid, North Kordofan State, Sudan, from January 2023 to December 2024. The trial involved a total of 224 participants. The trial either transported the patients to their homes or admitted them to the hospital during their labor, where they later gave birth. The sample offered an inclusive

portrayal of all the participants throughout the specified time period.

Statistician Analysis: We first organized the data into a data sheet and then input it into the Statistical Package for the Social Sciences (SPSS) (Version 24, Chicago, USA) computer software. We performed calculations for frequencies, percentages, cross-tabulation, and the chi square test. We derived the P-value from a 95% confidence interval (CI). Statistically significant results were defined as having p-values less than 0.05.

Informed consent: Before the interview, every participant had to sign a formal ethical consent form.

Ethical Approval: The esteemed Human Research Ethics Committee (HREC) at the Prof. Medical Research Center (MRCC) has duly accepted the protocol for this investigation. I hereby request that an approval number, MRCC 0010, be issued.

Results

This study investigated 224 women aged from 13 to 49 years, with a mean age of 28 years. Out of the 224 women, 68 (30.4%) were multipara, 99 (44.2%) were primigravida, and 57 (25.4%) were grandmultipara, as shown in Fig 1.




Figure 1 Provides a description of the study subjects by parity.

Most participants were aged ≤ 20 years, followed by 21–25 and ≥ 36 years, representing 50 (22.3%), 47 (21%), and 45 (20%), respectively. Out of the 244 women, 139 (62%) were from rural areas, and the remaining 85 (38%) were from urban areas. Fig. 3 shows the description of the study

subjects according to their rural or urban localities. The majority of participants were illiterate, followed by those with basic education level and university, constituting 67 (30%), 58 (26%), and 44 (20%), respectively, as indicated in Table 1, Fig 2.

Table 1. Distribution of the study subjects by parity and demographical characteristics

Variable	Multipara	Primigravida	Grandmultipara	Total
Age				
≤ 20 years	7	42	1	50
21-25	16	23	8	47
26-30	24	8	12	44
31-35	16	12	10	38
≥ 36	5	14	26	45
Total	68	99	57	224
Residence				
Rural	32	69	38	139
Urban	36	30	19	85
Total	68	99	57	224
Education				
Illiterate	7	41	19	67
Basic	23	20	15	58
Secondary	15	18	9	42
University	18	15	11	44
Postgraduate	5	5	3	13
Total	68	99	57	224




Figure 2. Description of the study subjects by demographic characteristics




Figure 3 Provides a description of the study population based on their localities.

Table 2, Figs. 4 and 5, summarize the distribution of research subjects based on parity and maternal outcomes. The most common maternal outcomes in this group of patients were postpartum hemorrhage, followed by puerperal sepsis and eclampsia, with 32 cases (14.3%), 29

cases (13%), and 20 cases (9%), respectively. We noticed variations for the different maternal outcomes when we counted the proportions with each parity group, as shown in Fig 5.

Table 2: Distribution of patients by parity and maternal outcomes.

Variable	Multipara	Primigravida	Grandmultipara	Total
Maternal outcomes				
Puerperal sepsis	7	15	7	29
Postpartum hemorrhage	8	11	13	32
Deep vein thrombosis	5	3	3	11
Eclampsia	3	15	2	20
Puerperal psychosis	4	9	2	15
Obstructed labour	5	4	7	16
Retained placenta	5	7	4	16
Birth canal injuries	4	3	2	9
Antepartum hemorrhage	4	12	4	20
No maternal complications	5	9	3	17
Rupture uterus	3	3	5	11
Vesicovaginal fistula	8	6	1	15
Death	7	2	4	13
Total	68	99	57	224




Figure 3. Description of the study subjects by Maternal outcomes




Figure 5. Maternal outcomes within the entire parity group

Table 3 and Fig 6 summarized the distribution of the study subjects by parity and fetal outcomes. Out of the 224 cases, only 69 (31%) were alive and well. Most cases were fresh

stillbirth, followed by macerated stillbirth, fetal distress, and birth trauma, which constituted 62 (28%), 38 (17%), 34 (15%), and 21 (9%), respectively.

Table 3: Patients are distributed by parity and fetal outcomes.

Variable	Multipara	Primigravida	Grandmultipara	Total
Fetal Outcomes				
Alive and well	23	30	16	69
Macerated still birth	10	16	12	38
Fresh stillbirth	22	24	16	62
Fetal distress	5	22	7	34
Birth trauma	8	7	6	21
Total	68	99	57	224




Figure 6. Proportions of fetal outcomes within entire parity groups

Fig. 7 displays the study participants' descriptions according to residence, mode of delivery, and birth attendant caliber. Most cases have spontaneous vaginal delivery followed by

Caesarean section and operative vaginal delivery, representing 87 (39%), 83 (37%), and 54 (24%), respectively. Fig. 7 illustrates the quality of birth attendants.




Figure 7 Provides a description of the study participants based on their residence, mode of delivery, and the caliber of their birth attendant.

Discussion

In April 2023, Sudan experienced a substantial conflict that had a profound impact on the healthcare system throughout the entire country. Maternal health and antenatal care are among the most seriously affected. Home birth was prevalent in Sudan, especially in rural regions, prior to the conflict. However, after the war, there was a significant increase in home births, which had negative implications. Therefore, this study aimed to assess the maternal and fetal outcomes of a group of women enrolled during the war.

In this particular group of patients, the predominant maternal outcomes observed were postpartum hemorrhage (14.3%), followed by puerperal sepsis (13%), and eclampsia (9%). Postpartum hemorrhage, a serious obstetric complication, is the leading and most preventable cause of maternal death during childbirth [7]. Globally, the major preventable cause of maternal mortality is postpartum hemorrhage, which is defined as a cumulative blood loss of 1,000 mL or more or blood loss accompanied by signs or symptoms of hypovolemia, regardless of delivery method. Among affluent countries, the United States has one of the highest maternal mortality rates, with postpartum hemorrhage accounting for around 14% of all maternal deaths [8].

Puerperal sepsis constitutes a substantial contributor to avoidable maternal mortality in both economically developed and developing nations. Sometimes, it ranks as the fourth or third most common direct cause of maternal mortality. Although puerperal sepsis is uncommon, it significantly contributes to maternal mortality, morbidity, and chronic complications, including secondary infertility [9]. Traditionally, the postpartum period refers to the six-week period following childbirth, during which infections are common and affect approximately 5 to 7% of women. Puerperal sepsis accounts for 10–15% of postpartum deaths and is one of the top five causes of maternal mortality worldwide. Infections are the primary cause of death following spontaneous or induced abortions. The medical burden resulting from these diseases is worsening due to the increasing bacterial resistance to commonly given antibiotics [9]. Factors such as the mother's level of education, the delivery method, and the presence or absence of pregnancy problems influence the incidence of puerperal sepsis. Ensuring thorough prenatal screenings, providing universal access to healthcare, implementing delivery procedures based on scientific evidence, and promoting education and awareness campaigns focused on the well-being of mothers are extremely important [10].

By expanding access to and promoting the early use of prenatal intervention, we can prevent eclampsia, a precursor to significant maternal morbidity [11].

The occurrence of preeclampsia is increasing worldwide, with emerging countries experiencing a higher rate of instability than developed countries. The World Health Organization (WHO) estimates that preeclampsia affects 2% to 10% of births worldwide [12, 13].

Following ten months of the crisis in Sudan, over 108,000 pregnant women are encountering substantial difficulties in accessing the essential medical aid they require. In conflict-affected regions, an alarming 67% of healthcare facilities are currently inaccessible, with some maternity hospitals being non-functional [14].

The study found that the most common fetal outcomes were fresh stillbirth (28%), followed by macerated stillbirth (17%), fetal distress (15%), and delivery trauma (9%). In spite of worldwide endeavors to diminish maternal and newborn mortality, stillbirths continue to pose a substantial public health obstacle in numerous low- and middle-income nations. District health systems, widely regarded as the foundation of health systems, play a crucial role in addressing the reported lack of data on stillbirths [15]. Labor induction has the potential to avoid stillbirth, and many accepted reasons for inducing labor prioritize this goal. Nevertheless, following childbirth, the child experiences escalating chances of mortality, frequently determined by gestational age. When deciding on the optimal timing for delivery, it is important to take into account the risk of postnatal death in the infant in order to minimize the risk of stillbirth [16].

Fetal intrauterine hypoxia induces fetal distress, posing a significant risk to both the fetus and the pregnant woman. The primary clinical technique for evaluating fetal distress is cardiotocography. Physicians exhibit variable interpretation of CTG data due to subjective variability, physicians interpret CTG data differently. Therefore, an alternate diagnostic technique for identifying fetal distress is necessary. The deep learning-based model for diagnosing fetal distress achieves high classification accuracy but requires a large number of parameters and processing resources, which presents difficulties in its practical application [17].

A targeted clinical evaluation can distinguish between birth traumas and congenital anomalies. Over the past few decades, the incidence of birth trauma has decreased steadily as a result of advancements in obstetrics and the greater utilization of cesarean delivery for cases of dystocia or challenging vaginal deliveries. The incidence of birth trauma decreased from 2.6 per 1,000 live births in 2004 to 1.9 in 2012. The use of forceps and vacuum-assisted deliveries has also declined over the past three decades [18]. According to research, negative interactions with healthcare professionals and maternity care system dysfunction significantly contribute to the occurrence of delivery trauma. Insufficient knowledge and support led women to experience a sense of helplessness and anguish as a result of making poor decisions. Insufficient support exacerbates women's encounters [19].

In conclusion, home birth is common in Sudan, although it is associated with negative consequences for both the mother and the baby. The predominant maternal outcomes are postpartum hemorrhage, followed by puerperal infection and eclampsia. The predominant fetal outcomes are fresh stillbirth, macerated stillbirth, fetal discomfort, and

birth trauma. It is highly advisable to prioritize the improvement of the healthcare system for maternity and antenatal care.

Acknowledgement

The authors express their gratitude to the individuals at El-Obeid, Obs & Gyn Hospital, for their assistance in gathering the data. We express our gratitude to the volunteers for generously dedicating their time and actively taking part.

Funding

This work has been funded by MRCC. Grant number: MRCC0010

Authors' contribution

Abdallah AAA: Conception, data collection, analysis, drafting, approval of the final version.

Mohamed NAA: Conception, data collection, approval of the final version.

Mohammed AA: Conception, data collection, approval of the final version.

K Nasralla: Conception, data collection, approval of the final version.

Ahmed HG: Conception, consultation, critical revision, approval of the final version.

Conflict of interest

Authors declare no conflict of interest

Data Availability

Data regarding this research can be requested from the corresponding authors.

References

- Boerma T, Campbell OMR, Amouzou A, Blumenberg C, Blencowe H, Moran A, Lawn JE, Ikilezi G. Maternal mortality, stillbirths, and neonatal mortality: a transition model based on analyses of 151 countries. *Lancet Glob Health*. 2023 Jul;11(7):e1024-e1031. doi: 10.1016/S2214-109X(23)00195-X.
- Chauke L, Bhoora S, Ngene NC. Postpartum haemorrhage - an insurmountable problem? *Case Rep Womens Health*. 2023 Feb 1;37: e00482. doi: 10.1016/j.crwh.2023.e00482.
- Ekwuazi EK, Chigbu CO, Ngene NC. Reducing maternal mortality in low- and middle-income countries. *Case Rep Womens Health*. 2023 Sep 9;39:e00542. doi: 10.1016/j.crwh.2023.e00542.
- University College Cork Ireland Definition and Classification of Maternal Death. 2023. <https://www.ucc.ie/en/mde/definitionandclassificationofmaternaldeath/>
- Perera SM, Isa GP, Sebushishe A, Sundararaj P, Piccirillo M, Xia S, Langaigane A, Ali J, Casey SE. "Midwives are heroes of the country": qualitative evaluation of a midwifery education program in South Sudan. *Front Glob Womens Health*. 2023 Aug 29; 4:1215405. doi: 10.3389/fgwh.2023.1215405.
- Abdelmola A. Antenatal Care Services in Sudan Before and During the 2023 War: A Review Article. *Cureus*. 2023 Dec 23;15(12):e51005. doi: 10.7759/cureus.51005.
- Turkoglu O, Friedman P. Evaluation During Postpartum Hemorrhage. *Clin Obstet Gynecol*. 2023 Jun 1;66(2):357-366. doi: 10.1097/GRF.0000000000000784.
- Jackson TL, Tuuli MG. Intrauterine Postpartum Hemorrhage-Control Devices. *Obstet Gynecol*. 2023 Nov 1;142(5):1000-1005. doi: 10.1097/AOG.0000000000005403.
- Sahle SG, Weldemariam S, Mehari MA, Abraha TA. Determinants of puerperal sepsis among post-partum mothers in Mekelle city public hospitals, Tigray, Ethiopia, 2021: a case control study. *BMC Womens Health*. 2023 Sep 21;23(1):502. doi: 10.1186/s12905-023-02643-2.
- Seboka K, Gurara AM, Bekele NT, Belachwe YA, Getahun MS, Negussie YM. Determinants of puerperal sepsis among postpartum women at a tertiary care hospital in Ethiopia: an unmatched case-control study. *Contracept Reprod Med*. 2024 Apr 24;9(1):18. doi: 10.1186/s40834-024-00283-x.
- Guglielminotti J, Daw JR, Friedman AM, Landau R, Li G. Medicaid expansion and risk of eclampsia. *Am J Obstet Gynecol MFM*. 2023 Aug;5(8):101054. doi: 10.1016/j.ajogmf.2023.101054.
- Osungbade KO, Ige OK. Public health perspectives of preeclampsia in developing countries: implication for health system strengthening. *J Pregnancy*. 2011;2011:481095. doi: 10.1155/2011/481095.
- Khan B, Allah Yar R, Khakwani AK, Karim S, Arslan Ali H. Preeclampsia Incidence and Its Maternal and Neonatal Outcomes with Associated Risk Factors. *Cureus*. 2022 Nov 6;14(11):e31143. doi: 10.7759/cureus.31143.
- Relief International. Pregnant Women and New Mothers Resilient Despite Ten Months of Conflict in Sudan. Accessed on 30/05/2024. Available at: <https://www.ri.org/pregnant-women-and-new-mothers-resilient-despite-ten-months-of-conflict-in-sudan/>
- Mensah Abrampah NA, Okwaraji YB, Oteng KF, Asiedu EK, Larsen-Reindorf R, Blencowe H, Jackson D. District health management and stillbirth recording and reporting: a qualitative study in the Ashanti Region of Ghana. *BMC Pregnancy Childbirth*. 2024 Jan 29;24(1):91. doi: 10.1186/s12884-024-06272-x.
- Rosenstein MG, Cheng YW, Snowden JM, Nicholson JM, Caughey AB. Risk of stillbirth and infant death stratified by gestational age. *Obstet Gynecol*. 2012 Jul;120(1):76-82. doi: 10.1097/AOG.0b013e31825bd286.
- Deng Y, Zhang Y, Zhou Z, Zhang X, Jiao P, Zhao Z. A lightweight fetal distress-assisted diagnosis model based on a cross-channel interactive attention mechanism. *Front Physiol*. 2023 Mar 6; 14:1090937. doi: 10.3389/fphys.2023.1090937.
- Dumpa V, Kamity R. Birth Trauma. 2023 Aug 28. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. PMID: 30969653.
- Watson K, White C, Hall H, Hewitt A. Women's experiences of birth trauma: A scoping review. *Women Birth*. 2021 Sep;34(5):417-424. doi: 10.1016/j.wombi.2020.09.