

Physiology-Driven Risk Stratification and Outcome Heterogeneity in Emergency Colorectal Cancer Presenting with Obstruction or Perforation: A Systematic Review

Abdel Latif Khalifa Elnaim^{1*}, Michael Pak Kai Wong², Ismail Sagap³

¹Department of Surgery Kassala Police Hospital Kassala -Sudan ORCID ID: 0000-0002-4763-6669 Email: almerfaby@gmail.com

²Sunway Medical Center Kuala Lumpur – Malaysia ORCID ID:0000-0001-9137-3096 Email: michaelpkwong@sunmed.my

³Department of Surgery, Faculty of Medicine, University Kebangsaan Malaysia (UKM), Hospital Canselor Tuanku Muhriz -UKM Kuala Lumpur ORCID ID: 0000-0003-2792-3529 Email: drisagap@gmail.com

| Article Type | Received | Accepted | DOI |
|------------------|---------------|-------------|----------------------------|
| Original Article | 27 April 2026 | 21 May 2026 | 10.70946/KJMHS3-2-26-P-101 |

ABSTRACT

Background: Obstructed and perforated colorectal cancer with emergency presentation represents a high-risk clinical condition associated with significant morbidity and mortality. Although outcomes are consistently inferior to elective surgery, marked variability was demonstrated in reported mortality, operative strategy, and long-term survival across studies leading to considerable heterogeneity. Differences in reporting physiological status, sepsis severity, tumor characteristics, and lack of standardized reporting contribute to this heterogeneity. **Objectives:** This systematic review aimed to assess current evidence on obstructed and perforated colorectal cancer, focusing on risk stratification, physiological determinants of outcome, and variability in operative frameworks and outcome reporting. **Methods:** A systematic search of electronic databases including PubMed, Embase, Scopus, Cochrane Library, and Google Scholar were performed to retrieve randomized trials, meta-analyses, cohort studies, and international guidelines addressing adult emergency colorectal cancer. Primary outcomes included postoperative morbidity and short-term mortality. Secondary outcomes included stoma formation, intensive care utilization, recurrence, and survival. The review findings were synthesized narratively in accordance with PRISMA guidelines because of substantial heterogeneity in definitions of emergency presentation, obstruction, perforation, and outcome measures. **Results:** Obstructed and perforated colorectal cancer presenting as emergency demonstrated consistently inferior outcomes compared to elective surgery; however, morbidity and mortality rates were variably reported across literature. Perforation was associated with the highest early mortality, particularly in patients presenting with sepsis and physiological instability. Operative strategies ranged from primary resection with anastomosis to staged surgery and Hartmann’s procedure, demonstrating variability in risk assessment and institutional expertise. Substantial variations in definitions and outcomes reporting limited direct inter-study comparability. **Conclusion:** Perforated and obstructed colorectal cancer presenting as emergency represent a high-risk surgical condition associated with considerable morbidity and mortality. Evidence generated in this review demonstrates that outcomes are primarily determined by physiological instability, septic burden, and system-level factors, rather than tumor characteristics alone. Perforation with generalized peritonitis carries the greatest early mortality, whereas obstructive disease characterized by a broader physiological spectrum in which risk is strongly modulated by baseline physiological reserve and the timeliness of resuscitative care.

Keywords: Colorectal cancer; emergency; obstruction; perforation; risk stratification; outcome heterogeneity.

Corresponding Author e-mail: almerfaby@gmail.com

INTRODUCTION

Introduce Colorectal cancer (CRC) is a major global health concern and among the most common malignancies worldwide, leading to substantial morbidity and mortality, with increasing incidence among younger population. Global cancer statistics database (GLOBOCAN 2020) estimate that CRC is one of the most commonly diagnosed malignancies with significant contributor to global cancer deaths [1,2,3,4]. A clinically significant proportion of patients continue to present as surgical emergencies, despite improvement in early detection and long survival in many reports secondary to population-based screening programs and advances in multimodal oncologic care [5,6].

Colonic obstruction or tumor perforation are the most common emergency presentations of CRC. Several studies demonstrated that emergency presentation is associated with poorer short and long-term outcomes compared to elective cohorts [7,8,9]. With variable screening uptake, socioeconomic class, definition of emergency, and referral pathway, reported rates of emergency presentation are considerably different between studies and healthcare systems [8,10,11]. Identified predeterminants of emergency presentation include socioeconomic disparities and delayed diagnostic pathways [11,12,13]. Although, obstruction and perforation both present as emergency, they represent pathophysiologically distinct clinical entities. Obstruction may present with minimal to maximal physiological instability along a spectrum from compensated colonic dilation to ischemia and impending perforation, while perforation is consistently accompanied by generalized peritonitis, systemic inflammatory response, and eventually organ failure [14,15,16]. Several studies demonstrated that perforation is associated with high early mortality and worse long-term oncologic outcomes, potentially acting as both a clinical and biological risk modifier [15,16,17,18]. These differences underscore the need to distinguish between subtypes of emergency presentation rather than treating them as a homogeneous entity.

Physiological status at emergency presentation is a predictor of outcomes in emergency CRC, beyond

tumour stage. Factors that increase postoperative morbidity and mortality include advanced age, comorbidity burden, and high American Society of Anesthesiologists (ASA) classification [19,20]. Additionally, postoperative complications in turn adversely affect long-term oncologic outcomes [21], suggesting that perioperative physiological instability may have durable prognostic implications. Importantly, emergency resection has been independently associated with reduced long-term survival even after adjustment for confounding factors [22]. Other factors related to system-level factors introduce additional differences. Institutional surgical volume, availability of surgical subspecialization, and perioperative protocols and resources impact short- and long-term outcomes in emergency CRC [23]. Variability in referral pathways and access to specialized service may partly explain inter-study heterogeneity in morbidity and mortality rates [7,11]. Consequently, interpretation of published data requires consideration of both patient-level and healthcare system-level determinants.

Surgical management in emergency CRC is inherently heterogeneous. Practical approaches include resection with primary anastomosis, Hartmann's procedure, staged diversion, and—in selected cases of left-sided obstruction—endoscopic decompression using self-expanding metal stents (SEMS) as a bridge to surgery [14,24-29]. Several reports from randomized and observational studies on SEMS show variable short- and long-term outcomes, and ongoing controversy persists regarding optimal patient selection [27,28,30-33]. Similarly, minimally invasive procedures and enhanced recovery protocols have been explored in selected emergency settings, but evidence remains heterogeneous and influenced by case selection [34-37].

Although, there is a growing body of literature focusing on emergency CRC management, substantial heterogeneity and variability persists in defining emergency presentation, classification of obstruction and perforation, reporting of mortality endpoints (in-hospital, 30-day, or 90-day), complication classification, and colonic diversion outcomes [38,39]. These differences constrain comparison among studies and complicates

synthesis of operative outcomes and survival data. Additionally, many comparative studies do not adjust for severity of physiological instability or sepsis status, potentially magnifying apparent variation between operative strategies.

This systematic review therefore aims to examine current evidence with particular focus on: (1) physiological determinants of outcome, (2) variability in operative decision-making, and (3) inconsistencies in outcome reporting standards, in order to clarify interpretation of existing data and identify priorities for methodological standardization in future research. Unlike several reviews focusing primarily on surgical procedures or stent placement, this review specifically evaluates emergency colorectal cancer through the lens of physiological risk stratification and methodological heterogeneity in outcome reporting. By focusing on physiological risk stratification and methodological heterogeneity, this review aims to provide a clinically relevant framework to improve interpretation of existing evidence and inform future research and guideline development in emergency colorectal cancer.

METHODS

Protocol and reporting framework:

This systematic review was reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. A written protocol explaining objectives, eligibility criteria, outcomes, and analysis protocol was finalized before study selection and was filed with the institutional research ethics board to document the planned methodology and enhance transparency. Two reviewers independently screened titles, abstracts, and full texts for eligibility. Disagreements were resolved through discussion, and where necessary by consultation with a third reviewer. The review protocol was finalized prior to study selection and approved by the institutional research ethics board; however, it was not prospectively registered in PROSPERO.

Search strategy and information sources

A systematic search was performed in electronic databases including PubMed, the Cochrane Library, Embase, Scopus, and Google Scholar to the most recent

search date. The search was directed to retrieve evidence relevant to adult colorectal cancer presenting as an emergency due to large bowel obstruction and/or perforation, including both management and outcome reporting. Search strategies combined controlled vocabulary (e.g., MeSH/Emtree terms where applicable) and free-text keywords including colorectal/colon/rectal cancer, emergency presentation, malignant large bowel obstruction, perforation, stoma/diversion, resection and anastomosis, Hartmann's procedure, self-expanding metal stents (SEMS), postoperative complications, mortality, recurrence, and survival. To maximize retrieval, the bibliographies of eligible articles and key reviews were hand-searched, and forward screening was performed where relevant to identify additional studies not captured in database queries. The full Boolean search strategy used for PubMed and adapted for other databases is provided in Appendix 1.

Eligibility criteria

Studies were considered eligible if they evaluated adult patients with colorectal cancer presenting emergently with obstruction and/or perforation and provided extractable information on outcomes and/or operative strategy. Eligible studies included randomized controlled trials (RCTs), observational cohort studies (prospective or retrospective), systematic reviews/meta-analyses, and international or national clinical guidelines.

Inclusion criteria

Studies were included when they met one or more of criteria such as reporting incidence or outcomes of emergency CRC presentations, evaluating postoperative morbidity and/or short-term mortality, reporting longer-term endpoints (recurrence, survival, or related oncologic outcomes), comparing emergency with elective CRC outcomes (when relevant to the review question), assessing emergency operative strategies (e.g., primary anastomosis ± diversion, Hartmann's procedure, staged diversion, SEMS as bridge to surgery), and/or addressing multidisciplinary care or perioperative pathway components applicable to emergency CRC (including modified ERAS elements).

Exclusion criteria

Excluded studies included case reports, letters, editorials, and non-systematic expert opinion pieces, studies

restricted to elective CRC populations without emergency data, studies focused on benign colorectal disease, pediatric-only cohorts, and non-English publications.

Study selection process

Articles identified through the search were collated, and screened after removal of duplicates. Screening of titles and abstracts against eligibility criteria, and eligible relevant records underwent full-text assessment. The selection process is illustrated in the PRISMA flow diagram (Figure 1). Using this strategy, 950 records were retrieved (850 from electronic databases and 100 from additional sources). After removal of duplicates, 480 records underwent title and abstract evaluation, of which 193 full-text articles were assessed for eligibility. Ultimately, 65 studies met the inclusion criteria and were included in the final qualitative synthesis after exclusion of pediatric cohorts, elective-only populations, case reports, and other ineligible study designs.




Figure 1 PRISMA 2020 Flow Diagram of Study Selection

RISMA 2020 flow diagram illustrating the study identification, screening, eligibility assessment, and inclusion process for this systematic review of emergency colorectal cancer presenting with obstruction or perforation.

Data extraction and variables of interest

Data were extracted using a structured template retrieving study setting and design, population characteristics, emergency type (obstruction, perforation, or mixed cohorts), operative strategy, and reported outcomes. Because the focus of this review includes risk stratification and outcome heterogeneity, additional extraction domains included the operational definitions used by authors for “emergency presentation” (e.g., admission route, timing to intervention), obstruction (clinical/radiologic/intraoperative criteria; complete vs partial when available), perforation (free vs contained; tumor vs diastatic when reported), mortality timepoints (in-hospital, 30-day, 90-day), complication reporting (overall vs graded systems), and stoma reporting (temporary diversion vs permanent end stoma where distinguishable). Primary outcomes were postoperative morbidity and short-term mortality (as defined by each study). Secondary outcomes included stoma formation, length of hospital stay, intensive care utilization, recurrence, and survival endpoints. Data extraction was performed independently by two reviewers using a standardized template.

Quality assessment

Study quality was appraised using design-appropriate tools such as AMSTAR 2 for systematic reviews/meta-analyses, the Cochrane Risk of Bias tool for RCTs, the Newcastle–Ottawa Scale for cohort studies, AGREE II for clinical guidelines, and SANRA for narrative/scoping reviews. A structured summary of appraisal findings is provided in Appendix 2.

Data synthesis

Taking into consideration the substantial heterogeneity across included articles formal meta-analysis was not performed, results were presented narratively with structured tables underscoring differences between obstructed and perforated disease, the influence of physiological risk and case-mix on outcomes, and variability in reporting standards that limits inter-study comparability. Summary statistics were reported where clinically appropriate and methodologically relevant, without combining effect estimates across incompatible definitions.

Ethical considerations

This systematic review is not presenting any identifiable patients' information and not applying any intervention. The included articles had passed their ethical approval therefore; additional ethical approval was not required.

RESULTS

Study selection and evidence profile

The included studies were methodologically diverse, comprising registry-based analyses, retrospective and prospective cohort studies, randomized trials, systematic reviews, and international guideline statements

[14,25,40,41]. Geographical distribution of included studies covered Europe, the Americas, Asia, the Middle East, and Africa, reflecting considerable geographic variability in healthcare infrastructure and screening programs [8,11, 23]. Across studies, heterogeneity was evident at multiple levels including case definitions, phenotyping of obstruction and perforation, physiological instability reporting, operative approaches, and outcome measurement. This variability informed the decision to prioritize structured narrative synthesis rather than quantitative pooling. (Table 1)

Table 1 Characteristics of Studies Included in the Systematic Review and Primary Sources of Methodological Heterogeneity

| Study (Author, Year) | Country | Study Design | Study Population | Emergency Phenotype | Primary Outcomes Reported | Key Methodological Features Influencing Heterogeneity |
|-------------------------|--------------|-------------------------------------|----------------------------------|---------------------------|--|---|
| Pisano et al., 2018 | Italy | International guideline (WSES) | CRC with obstruction/perforation | Obstruction & perforation | Mortality, morbidity, operative strategy | Guideline-based consensus; heterogeneous source evidence; no uniform physiological stratification |
| Ansaloni et al., 2010 | Italy | Consensus guideline | Left-sided obstructing CRC | Obstruction | Operative options, short-term outcomes | Focused on left-sided disease; limited long-term oncologic reporting |
| Golder et al., 2022 | UK | Systematic review & meta-analysis | CRC emergency presentation | Mixed emergency CRC | Incidence, short-term mortality | Administrative coding variability; mixed mortality timepoints |
| Ma W et al., 2023 | China | Systematic review & meta-analysis | Obstructive CRC | Obstruction | Mortality, complications, SEMS outcomes | Selection bias in SEMS cohorts; exclusion of perforated/septic patients |
| Esswein et al., 2023 | Austria | Matched retrospective cohort | Emergency CRC resection | Emergency CRC | Long-term survival, morbidity | Case-mix severity adjustment; residual confounding by indication |
| Arnarson et al., 2023 | Sweden | Registry-based retrospective cohort | Emergency colon cancer | Emergency CRC | Mortality, overall survival | Subspecialization effect; institutional volume influence |
| Kaewubon et al., 2024 | Thailand | Retrospective cohort | Stage II emergency colon cancer | Emergency CRC | Recurrence, prognostic factors | Stage-restricted cohort; perforation as recurrence modifier |
| Chen et al., 2017 | Taiwan | Retrospective cohort | CRC with obstruction/perforation | Obstruction & perforation | Mortality, short-term outcomes | Variable perforation definition (free vs contained not standardized) |
| Moura et al., 2014 | Brazil | Retrospective cohort | Emergency CRC surgery | Emergency CRC | Stoma rate, mortality | High stoma prevalence; unclear distinction between temporary vs permanent |
| Ocak et al., 2021 | Turkey | Retrospective cohort | Emergency oncologic CRC | Emergency CRC | Morbidity, mortality | Mortality timepoint not uniformly specified |
| Bin Traiki et al., 2023 | Saudi Arabia | Retrospective cohort | Emergency CRC surgery | Emergency CRC | Outcomes, survival | Regional referral bias; mixed colon/rectal denominators |

| | | | | | | |
|------------------------|--------------|-------------------------------------|---------------------------|---------------------------|----------------------|--|
| Öistämö et al., 2016 | Sweden | Retrospective cohort | Emergency CRC | Emergency CRC | Mortality | Limited physiological severity reporting |
| Krutsri et al., 2021 | Thailand | Retrospective cohort | Elderly emergency CRC | Emergency CRC | Morbidity, mortality | Age-stratified cohort; high comorbidity burden |
| Kobayashi et al., 2023 | Japan | Multi-institutional registry cohort | Obstructed/perforated CRC | Obstruction & perforation | Mortality | Registry-based design; limited contamination grading |
| Norman et al., 2023 | South Africa | Retrospective cohort | Emergency CRC | Emergency CRC | Mortality, survival | Resource-limited setting; system-level variability |

- CRC = colorectal cancer.
- NR = not reported or not extractable from published data.
- Study populations varied in inclusion of colon-only versus combined colon and rectal cancers.

Variability in definitions of emergency presentation

Across studies, operational definitions of “emergency presentation” varied substantially. Some studies defined emergency status based on acute admission requiring urgent surgery, whereas others relied on administrative coding or timing from admission to operation [7,8,11]. Few studies clearly differentiate between emergency admission and emergency operative intervention. Emergency presentation was frequently linked to socioeconomic variations and delayed diagnostic pathways [8,11,13]. However, the threshold at which a presentation was labeled “emergent” was inconsistently reported, limiting cross-study comparability.

Heterogeneity in phenotyping obstruction and perforation

Considerable variation was identified in definitions of obstruction and perforation. Obstruction was variably classified using clinical criteria (e.g., obstipation, abdominal distension), radiologic findings (CT-confirmed transition point), or intraoperative confirmation [38,42,43]. Few studies distinguished complete from partial obstruction [42]. While perforation definitions ranged from free intraperitoneal perforation to contained or tumor-related

microperforation; differentiation between diastatic and tumor perforation was inconsistently specified [15,16,18,44]. Incidence estimates therefore varied widely as demonstrated in Table 2. Emergency presentation overall was frequently reported in the 15–45% range [2,3,40], but obstruction and perforation proportions differed markedly across cohorts [15,16,42]. These variable definitions likely contributed significantly to the broad reported ranges.

Physiological instability and risk stratification reporting

Reporting of baseline physiological status was inconsistent across studies. Worse postoperative outcomes were observed with advanced age, comorbidity burden, and higher ASA class [19,20]. However, detailed documentation of hemodynamic instability, sepsis severity, organ failure, or contamination grading was absent in many reports. Where physiological variables were reported, they identified strong associations with short-term mortality, particularly in perforated colonic tumors with generalized peritonitis [15,16,18]. Nonetheless, absence of standardized severity metrics limited relevant comparison between operative strategies and between obstructed and perforated cohorts.

Table 2 Incidence of Emergency Presentation, Obstruction, and Perforation in Colorectal Cancer: Impact of Definition and Mortality Timepoint Variability

| Study (Country, Year) | Study Design | Definition of Emergency Used | Emergency CRC (%) | Obstruction (%) | Perforation (%) | Mortality Timepoint Reported |
|--------------------------|----------------------|------------------------------------|-------------------|-----------------|-----------------|------------------------------|
| Kaewubon 2024 (Thailand) | Retrospective cohort | Urgent admission requiring surgery | 37.3 | NR | 5.5 | 30-day |
| Esswein 2023 (Austria) | Matched cohort | Emergency operative resection | 30.0 | NR | 2–9 | 90-day |
| Storli 2023 (Norway) | Registry analysis | Acute presentation | 11.0 | NR | NR | NR |
| Bin Traiki 2023 (KSA) | Retrospective cohort | Emergency surgery | 32.5 | 8.4 | 12.9 | In-hospital |
| Nahar 2022 (KSA) | Cohort | Admission-based | 33.0 | NR | NR | NR |
| Maertens 2022 (UK) | Case series | Emergency surgical episode | 25.0 | NR | NR | NR |
| Golder 2022 (UK) | Systematic review | Administrative coding | 20.0 | 14.4 | NR | Mixed |
| Ocak 2021 (Turkey) | Retrospective | Emergency resection | 18.0 | NR | NR | NR |
| Acar 2020 (Turkey) | Retrospective | Urgent oncologic surgery | 18.4 | NR | NR | In-hospital |
| Pisano 2018 (WSES) | Guideline | Clinical emergency definition | 13.2 | 3.4 | NR | Not standardized |

- Percentages refer to the proportion reported within each individual study population. Mortality percentages reflect the timepoint specified in each original study (in-hospital, 30-day, or 90-day).
- Definitions of “emergency presentation,” “obstruction,” and “perforation” varied across studies and were based on authors’ criteria (e.g., clinical presentation vs operative findings).
- NR = not reported in the original publication.

Short-term outcomes

Across studies, emergency CRC was consistently associated with worse short-term outcomes compared to elective surgery [7,9,22]. However, reported mortality rates varied substantially, influenced by case-mix, institutional capacity, and mortality timepoint definitions (in-hospital vs 30-day vs 90-day) as demonstrated in **Table 3**. Perforated CRC demonstrated the highest early mortality in most studies [15,16,18], whereas obstructed CRC outcomes were more heterogeneous and appeared strongly modulated by physiological reserve [19,29]. Postoperative morbidities were independently associated with adverse long-term oncologic outcomes in several studies [21], suggesting that perioperative physiological insult may extend beyond short-term morbidity.

Resource utilization and perioperative burden

Length of hospital stay (LOS) and ICU utilization were frequently reported but inconsistently defined. Some

studies revealed mean LOS, others median values; and criteria for ICU admission were rarely standardized. Higher ICU use and prolonged hospitalization were most often observed in patients with sepsis, hemodynamic instability, or multi-organ dysfunction [15,16,20]. Institutional variations in perioperative pathways and system-level resources likely contributed additional variability [23,45].

Long-term outcomes

Long-term endpoints showed considerable heterogeneity. Overall survival (OS), disease-free survival (DFS), cancer-specific survival (CSS), and recurrence rates were variably reported. Follow-up duration ranged widely across studies, and censoring practices were inconsistently described [39,46, 47,48]. Emergency surgery was associated with inferior long-term survival in several matched or registry-based analyses, even after adjustment for confounders [22,49,50,51]. Perforation

was a particularly adverse prognostic factor, including in stage II disease [51]. However, inconsistent endpoint

definitions and surveillance intervals limited quantitative synthesis and prevented robust comparative conclusions.

Table 3 Short- and Long-Term Outcomes of Emergency Colorectal Cancer Surgery: Impact of Mortality Timepoint, Follow-Up Duration, and Cohort Phenotype

| Study (Country, Year) | Emergency Cohort Phenotype | Stoma Formation (%) | Morbidity (%) | Mortality (%) | Mortality Time point | Recurrence (%) | Survival Endpoint Reported | Follow-up Duration | Key Source of Outcome Variability |
|-----------------------------|--|---------------------|---------------|---------------|----------------------|----------------|----------------------------|--------------------|---|
| Moura (Brazil, 2014) | Mixed emergency CRC | 85.0 | 33.4 | 50.0 | In-hospital | NR | NR | NR | High permanent stoma rate; unclear stoma classification |
| Ma W (China, 2023) | Obstructive CRC | NR | 53.4 | 50.0 | 30-day | NR | OS (77.8%) | Variable | SEMS selection bias; obstruction-only cohort |
| Elmessiry (Egypt, 2023) | Curative emergency CRC | NR | 11.1 | 28.9 | 30-day | 25.2 | OS (71.1%) | 3–5 years | Curative-only selection; lower physiological severity |
| Arnarson (Sweden, 2023) | Emergency colon; subspecialist vs generalist | 42.5 | 35.4 | 33.4 | 90-day | 28.3 | OS (47.0%) | Long-term | Volume and subspecialization effect |
| Boeding (Netherlands, 2023) | Right-sided obstructing CRC | 29.0 | 10.0 | NR | Not specified | NR | OS (60.9%) | Long-term | Right-sided disease focus; registry data |
| Zamaray (Netherlands, 2023) | Emergency CRC | NR | 29.0 | NR | Not specified | NR | NR | NR | Mortality endpoint not uniformly defined |
| Pisano (UK, 2018) | CRC emergencies (guideline synthesis) | NR | NR | 25.2 | Mixed | NR | NR | NR | Heterogeneous source data |
| Öistämö (Sweden, 2016) | Emergency CRC | 6.5 | 26.0 | 12.0 | 30-day | NR | NR | NR | Registry-based; limited physiological data |
| Esswein (Austria, 2023) | Emergency resection | 39.0 | 47.4 | 47.5 | 90-day | 35.6 | NR | Long-term | Matched analysis; residual confounding |
| Krutsri (Thailand, 2021) | Elderly emergency CRC | NR | 41.3 | 26.3 | In-hospital | NR | NR | NR | Age-stratified high-comorbidity cohort |
| Kobayashi (Japan, 2023) | Obstruction & perforation | 8.3 | 18.5 | 53.3 | In-hospital | NR | NR | NR | Mixed obstruction/perforation; contamination grade unreported |
| Norman (South Africa, 2023) | Emergency CRC | 4.3 | 31.0 | 2.2 | 30-day | 5.4 | OS (55.4%) | Variable | Resource-limited setting; referral bias |

| | | | | | | | | | |
|------------------------|---------------|------|------|------|---------------|----|------------|----------|---|
| Bin Traiki (KSA, 2023) | Emergency CRC | 28.6 | NR | NR | Not specified | NR | OS (60.9%) | Variable | Regional tertiary referral cohort |
| Ocak (Turkey, 2021) | Emergency CRC | NR | 31.0 | NR | Not specified | NR | NR | NR | Mortality timepoint unspecified |
| Kim (Korea, 2010) | Emergency CRC | NR | NR | 45.9 | In-hospital | NR | NR | NR | Early-era cohort; unspecified severity stratification |

- Mortality percentages are not directly comparable across studies due to differences in reporting timepoints and baseline physiological severity.
- Survival percentages reflect study-specific follow-up durations and should be interpreted within individual study context.
- Mortality endpoints varied across studies (in-hospital, 30-day, or short-term postoperative mortality).
- Morbidity definitions were heterogeneous and included overall complications or major complications as defined by individual authors.
- Survival outcomes represent overall survival where reported; follow-up duration varied between studies.
- NR = not reported.

Operative strategy variation

As illustrated in **Table 4** operative approaches included primary resection with anastomosis (with or without diversion), Hartmann’s procedure, staged diversion, and SEMS as a bridge to surgery in selected left-sided colonic obstruction [14,25,27,28,29,32,33,52]. Operative strategies differed considerably across studies and healthcare systems. In several reports, higher-risk or septic patients were more likely to undergo diversion or Hartmann’s procedure, reflecting confounding by indication [19,22]. Current multidisciplinary reviews highlight the importance of structured collaboration between colorectal surgeons, endoscopists,

anesthesiologists, and oncologists in the management of malignant colonic obstruction, particularly on performing SEMS placement versus immediate resection [53]. SEMS outcomes differed across randomized and multicenter trials, with variations in technical success, perforation risk, and oncologic endpoints [27,28,32,33,54]. Guideline recommendations consistently emphasized selective application based on physiological stability and institutional expertise [14,25,41] (Figure 2). Minimally invasive approaches were reported in specialized settings but were influenced by case selection, surgeons’ experience, and institutional capacity [34,55].

Table 4 Risk-Stratified Emergency Management Strategies in Obstructed or Perforated Colorectal Cancer: Evidence Strength, Physiological Context, and Sources of Controversy

| Clinical Scenario | Strategy / Approach | Evidence Level in Included Literature | Physiological Context Driving Selection | Principal Controversy | Structured Practical Implication |
|----------------------------|--|--|---|---|--|
| Obstructing CRC (general) | Immediate emergency resection (various procedures) | Observational cohorts; registry analyses | Variable physiological reserve; risk of ischemia or impending perforation | High morbidity and mortality; outcomes strongly influenced by case-mix and institutional capacity | Prioritize preoperative resuscitation; document physiological status; centralize to experienced colorectal units when feasible |
| Left-sided obstructing CRC | SEMS as bridge-to-surgery (BTS) | RCTs, meta-analyses, guideline updates | Hemodynamically stable; absence of perforation or generalized peritonitis | Perforation risk; potential oncologic compromise; selection bias in stable patients | Use selectively in stable patients; explicitly report technical success, perforation rates, and selection criteria |

| | | | | | |
|------------------------------------|--|--|--|--|---|
| Left-sided obstruction | Diverting stoma / staged surgery | Guidelines; consensus statements; cohort studies | Physiological instability; high ASA; uncertain anastomotic safety | Balancing delayed definitive resection vs reduced leak risk | Recommended in unstable or high-risk patients; include rationale (physiology vs contamination) in operative documentation |
| Obstructing CRC | Primary resection + anastomosis (± diversion) | Guidelines; observational cohorts | Hemodynamically stable; limited contamination; adequate tissue perfusion | Anastomotic leak risk vs oncologic adequacy; confounding by indication | Restrict to carefully selected stable patients; require explicit intraoperative risk assessment documentation |
| Perforated CRC (localized or free) | Damage-control principles / staged reconstruction | Guidelines; retrospective perforation cohorts | Septic physiology; generalized peritonitis; organ dysfunction | One-stage reconstruction vs staged management | Source control and sepsis resuscitation take precedence; consider staged reconstruction after stabilization |
| Perforated CRC | Hartmann's procedure / resection without anastomosis | Cohort studies; guideline recommendations | Shock, fecal peritonitis, severe contamination, high ASA | Reduced leak risk vs permanent stoma burden | Define institutional criteria for Hartmann; mandate reporting of permanent vs temporary stoma outcomes |
| Perforated CRC (stage II subgroup) | Recurrence risk stratification | Observational stage-specific cohorts | Perforation as biological and inflammatory modifier | Whether perforation mandates intensified adjuvant therapy | Incorporate perforation status into multidisciplinary oncologic discussion and surveillance planning |
| Operator-related factor | Specialist colorectal vs general surgeon | Registry-based cohort studies | Institutional volume; subspecialization availability | Volume-outcome relationship; access disparities | Establish referral networks and on-call colorectal pathways; audit outcomes stratified by surgeon specialty |
| Surgical approach | Minimally invasive (laparoscopy/robotics) | Case series; selected cohorts | Hemodynamic stability; limited contamination | Feasibility vs oncologic adequacy under emergency conditions | Restrict to stable patients with institutional expertise; avoid compromise of oncologic principles |
| Perioperative pathway | Adapted ERAS in emergency CRC | Systematic reviews; prospective cohorts | After initial resuscitation; stabilized physiology | Applicability in unstable or septic patients | Implement pragmatic emergency-ERAS bundle; clearly distinguish from elective ERAS protocols |

- This table summarizes *management-relevant contributions* of included sources; effect sizes and direct head-to-head comparisons were not consistently extractable across studies.
- Evidence level reflects study design within included literature and does not represent formal GRADE classification
- “NR” indicates details were not reported in the extracted dataset or were not comparable across studies.
- CRC = colorectal cancer; SEMS = self-expanding metal stent; BTS = bridge to surgery; ERAS = enhanced recovery after surgery.




Figure 2 . Evidence-Based Management Algorithm for Emergency Obstructed or Perforated Colorectal Cancer

Evidence-based decision pathway for the management of colorectal cancer presenting as an emergency with acute obstruction or perforation. Algorithm synthesized from included guideline statements and cohort evidence; intended as a conceptual risk-stratified framework rather than prescriptive protocol.

Reporting inconsistencies as a source of outcome heterogeneity

A central finding across included studies was the absence of harmonized reporting standards. Variability was observed in definition of emergency presentation [7,8], classification of obstruction and perforation [15,38], mortality timepoints (in-hospital vs 30-day vs 90-day), complication classification systems, reporting of temporary versus permanent stoma, and long-term oncologic outcomes [39,47]. This methodological heterogeneity constrained direct inter-study comparison and precluded meta-analysis (Figure 3). Variations in definitions and baseline severity reporting likely account

for a substantial proportion of the variability observed in documented morbidity and mortality across the studies. These findings highlight that heterogeneity in methodology and documentation contributes considerably to apparent differences in outcomes across published studies. (Table 5)

Summary of key results

Across 65 studies, outcome variability in emergency obstructed or perforated CRC appears strongly influenced by inconsistent definitions of emergency presentation and disease subtype, variable reporting of physiological severity and contamination, confounding by indication in operative strategy selection, divergent mortality timepoints and endpoint definitions, and differences in institutional infrastructure and system-level capacity.




Figure 3 Sources of Outcome Heterogeneity in Emergency Colorectal Cancer: Implications for Evidence Interpretation

Conceptual illustration of major sources of heterogeneity influencing outcomes and interpretation of evidence in emergency colorectal cancer.

Table 5 Proposed Harmonized Reporting Framework for Emergency Obstructed or Perforated Colorectal Cancer: Addressing Definition Variability and Methodological Heterogeneity

| Domain | Definition Applied in This Review | Observed Variability Across Studies | Consequence for Evidence Synthesis | Analytical Handling in This Review | Recommended Standardized Reporting Framework |
|--------------------------------------|--|--|--|--|--|
| Emergency presentation | Unplanned admission requiring urgent intervention for CRC-related obstruction or perforation | Defined variably by admission route, operative timing, administrative coding, or unclear criteria | Distorts reported incidence; impairs inter-study comparability | Incidence reported per study without pooling across inconsistent definitions | Emergency CRC defined as unplanned admission requiring operative or interventional management within 24–72 hours due to obstruction or perforation |
| Obstruction | Clinical and/or radiologic large bowel obstruction attributable to CRC | Complete vs partial obstruction inconsistently specified; CT vs clinical vs intraoperative diagnosis; inclusion of ileus | Influences operative selection (SEMS vs diversion vs resection); confounds outcome comparison | Obstruction rates retained as published; contextualized narratively | CT-confirmed mechanical obstruction secondary to CRC, stratified as complete vs partial, with documentation of ischemia/perforation risk |
| Perforation | CRC-associated perforation at presentation or during index surgery | Free vs contained perforation inconsistently distinguished; tumor vs diastatic perforation often merged; sepsis status variably reported | Major determinant of mortality and operative strategy; pooling unreliable without subtype stratification | Interpreted as marker of septic physiological burden | Mandatory classification: (1) tumor perforation, (2) diastatic perforation, (3) free vs contained, with documented sepsis severity and contamination grade |
| Baseline physiological status | ASA class and comorbidity data where available | Hemodynamic instability, organ dysfunction, and contamination grading inconsistently reported | Limits risk adjustment; exaggerates apparent operative strategy differences | Interpreted qualitatively in narrative discussion | Mandatory reporting of: ASA class, shock status, vasopressor use, organ dysfunction, and contamination severity |
| Mortality | Short-term postoperative mortality as reported by authors | In-hospital vs 30-day vs 90-day mortality inconsistently reported | Prevents valid comparison; inflates perceived heterogeneity | Mortality retained as reported; timepoint variability explicitly identified | Mandatory reporting of 30-day mortality; optional 90-day mortality; specify in-hospital deaths separately |
| Morbidity / complications | Postoperative complications following emergency CRC surgery | Overall vs major complications; inconsistent grading systems | Restricts severity stratification and benchmarking | Not pooled; interpreted descriptively | Complications graded using Clavien–Dindo classification; specify major morbidity (\geq III) |
| Stoma formation | Any stoma created during index admission | Temporary vs permanent stoma often not distinguished; | Overestimates permanent stoma burden; misrepresents | Reported as given; limitations acknowledged | Mandatory distinction between temporary diverting stoma and permanent end stoma; |

| | | | | | |
|---|--|---|--|---|--|
| | | reversal rates rarely reported | long-term impact | | report reversal rate and follow-up duration |
| Recurrence | Any reported oncologic recurrence post-emergency surgery | Local vs distant recurrence inconsistently separated; follow-up duration variable; stage-specific cohorts | Survival bias; non-comparable recurrence estimates | Reported only when explicitly defined | Report local and distant recurrence separately; minimum follow-up 3–5 years; specify stage distribution |
| Survival | Overall survival (OS) where available | OS vs DFS vs cancer-specific survival; heterogeneous follow-up intervals | Precludes pooled survival analysis | Presented descriptively without meta-analysis | Primary endpoint: OS; secondary: DFS; define fixed follow-up intervals (3- and 5-year) |
| Population denominator | Emergency CRC presenting with obstruction and/or perforation | Colon-only vs colon+rectum; resected-only vs all admissions; guideline vs registry populations | Denominator shifts distort incidence and mortality rates | Populations clearly labeled per study | Explicit reporting of denominator (all emergency admissions vs resected cases); specify colon vs rectal cancer inclusion |
| Operative strategy documentation | Procedure type as reported | Rationale for diversion vs anastomosis rarely documented; confounding by indication unmeasured | Limits causal interpretation of strategy comparisons | Interpreted within confounding framework | Mandatory documentation of operative rationale (physiology, contamination, technical factors) |

- Proposed definitions represent authors’ synthesis based on observed reporting variability and are not derived from formal consensus methodology.
- Recommendations are intended to guide future observational research and registry design rather than replace existing international guideline criteria.
- CRC = colorectal cancer; NR = not reported.
- Substantial heterogeneity in definitions and outcome reporting justified a structured narrative synthesis rather than quantitative pooling.
- The proposed standardized definitions are intended to inform future observational studies, registries, and guideline development.

DISCUSSION

1. Emergency CRC as a physiology-dominated clinical state

The collective evidence synthesized in this review demonstrates that emergency colorectal cancer (CRC) presenting with obstruction or perforation should be considered primarily as a physiologically unstable oncologic state, rather than simply an alternative presentation of tumor location or stage. Across several studies, short-term mortality is consistently higher in emergency resections compared with elective surgery [7,8,9,22]. However, this increased risk is not uniformly distributed and appears strongly influenced by baseline

physiological status. Perforation with generalized peritonitis and systemic inflammatory response, is consistently associated with the highest early mortality [15,16,18]. In contrast, obstruction demonstrates a broader risk spectrum, ranging from relatively compensated dilation to impending ischemia or septic deterioration [15,56]. The physiological reserve is identified as the predominant factor influencing the postoperative risk as demonstrated in studies that adjust for age, comorbidity burden, and ASA classification [19,20]. Comparative evaluations of perioperative risk prediction models in major abdominal surgery further revealed variability in predictive accuracy between ASA, POSSUM-derived systems, and NSQIP-based tools, highlighting the need for standardized risk adjustment in

emergency CRC research [57]. Additionally, postoperative morbidities have been independently linked to worse long-term oncologic outcomes [21]. This demonstrates that perioperative physiological stress may not merely influence short-term morbidity but may also alter long-term disease pathways. These findings support a strategy in which physiological instability and sepsis severity act as primary determinants of outcome, with tumor factors contributing secondarily.

2. Confounding by indication and operative strategy interpretation

Across the studies, a recurrent limitation is confounding by indication. Operative strategy in emergency CRC is guided and influenced by intraoperative judgment, physiological stability status, sepsis burden, and institutional expertise. Higher-risk patients presenting with septic shock, diffuse peritonitis, or limited physiological reserve are more likely to undergo diversion or Hartmann's procedure [14,41], whereas primary anastomosis is preferentially performed in more stable patients [57,58]. Consequently, comparative analyses between approaches often reflect baseline case-mix variations rather than intrinsic superiority of one strategy. Similarly, the literature on self-expanding metal stents (SEMS) demonstrates how selection bias influences outcomes. Randomized trials and multicenter studies (Dutch Stent-In, ESCO, CReST) demonstrate variable short- and long-term findings [27,28,32,33], but eligibility criteria typically exclude patients with overt perforation or severe peritonitis. Observed variations therefore suggest patient selection and procedural expertise rather than a uniform therapeutic effect [25,30]. Economic analyses further demonstrate that the cost-effectiveness of SEMS as a bridge to surgery is highly context-dependent, influenced by institutional expertise, morbidity rates, and reduce stoma reversal probabilities [59]. Minimally invasive procedures were reported in specialized settings but were influenced by case selection and institutional volume and expertise [34,54]. Recent reviews reveal that minimally invasive emergency colorectal surgery remains largely restricted to hemodynamically stable patients managed in high-volume centers, reinforcing the central role of physiological selection in interpreting outcomes [60].

Registry-based analyses show that emergency resection is associated with worse long-term survival even after adjustment for confounders [4,19], but the extent to which residual physiological confounding persists is unclear. Without structured standardized documentation of sepsis grade, shock parameters, and organ failure, interpretation of operative comparisons remains inherently limited.

3. Reporting heterogeneity as a structural limitation

A major observation of this review is the lack of harmonized reporting standards across studies, resulting in substantial methodological heterogeneity. Variability was demonstrated in definitions of emergency presentation [7,8,11], classification of obstruction (clinical vs radiologic vs intraoperative) [38,42], perforation subtyping (free vs contained; tumor vs diastatic) [15,16,18], mortality timepoints (in-hospital, 30-day, 90-day), complication classification systems, reporting of temporary versus permanent stoma, and long-term oncologic endpoints such as overall survival (OS), disease free survival (DFS), and cancer specific survival (CSS) [39,47,48]. These inconsistencies magnify apparent inter-study differences and constrain quantitative synthesis. For example, mortality comparisons are inherently unreliable as some studies report in-hospital mortality while others report 90-day mortality. Similarly, recurrence patterns cannot have relevant comparison when follow-up intervals and outcome ascertainment methods vary. Importantly, only a limited number of studies stratify outcomes by physiological severity or contamination grade, despite these factors emerging as major determinants of risk. The absence of this stratification constrains attempts to generate practice-defining conclusions from pooled observational data.

4. Toward a minimum reporting dataset for emergency CRC

Based on the heterogeneity revealed, this review emphasizes the need for a structured standardized minimum reporting dataset for emergency obstructed and perforated CRC which should include, at minimum:

I. Baseline physiological variables

These include age and comorbidity indices, ASA classification, hemodynamic parameters such as shock status, vasopressor requirement, sepsis classification, organ failure markers, and frailty assessment in older adults using validated frailty indices, given contemporary evidence that frailty is an independent predictor of postoperative morbidity and mortality in colorectal cancer surgery [61]

II. Disease phenotype classification

This includes obstruction if it is complete or partial; radiologic criteria specified [38,42], perforation (free vs contained; tumor vs diastatic) [15,18], and contamination grading (localized vs diffuse peritonitis).

III. Operative documentation

The data should include procedure type such as primary anastomosis ± diversion, Hartmann's, staged procedure, justification for diversion such as physiology, contamination, or technical factors, and SEMS technical success and complications where applicable [27,28,32,33].

IV. Outcome definitions

This demonstrates that mortality reported at standardized intervals (30-day and/or 90-day), morbidities graded using a recognized system, clear distinction between temporary and permanent stoma, and consistent definition of long-term oncologic endpoints (OS, DFS, CSS) [39,47]. Adoption of such a standardized dataset would resolve interpretive ambiguity and improve comparability across institutions and studies.

5. Implications for guidelines and registry-based research

Existing international guidelines appropriately emphasizes tailored management based on physiological stability and contamination severity [14,25,41]. However, recent data from resource-limited settings show that comparative outcomes between bridge-to-surgery strategies and immediate emergency resection are strongly influenced by institutional volume and expertise and perioperative support capacity, further underscoring system-level determinants of outcome heterogeneity [62]. Large-scale registry initiatives have revealed the capacity to generate more robust outcome

analyses when standardized variables are collected [63,64]. Expansion of such registry approach incorporate physiological stratification and contamination grading could considerably enhance benchmarking and evidence synthesis. Additionally, incorporation of standardized emergency CRC modules within national colorectal registries may allow risk-adjusted outcome comparison across centers, evaluation of subspecialized service and institutional capacity effects [23] (Figure 4), and assessment of perioperative trajectories implementation [35,36,45]. Future registry-linked assessments may also evaluate perioperative systemic therapy protocols in selected emergency presentations, including neoadjuvant therapy in biologically advanced but operable colon cancer [65]. Similar system-level standardization may reduce variability and clarify the true comparative effectiveness of operative strategies.

The findings of this review demonstrate that emergency colorectal cancer should be formulated as a physiologically-driven surgical condition characterized by varying degrees of septic burden and physiological instability and not merely as an oncologic presentation. Understanding of outcomes across literature is therefore strongly influenced by variations in baseline physiological derangement, sepsis status, and institutional infrastructures rather than tumor characteristics alone. Recognizing emergency colorectal cancer as a physiology-dominated condition may facilitate improved risk stratification, more relevant interpretation of operative outcomes, and more reliable comparisons across international datasets.

Strengths: This systematic review integrates evidence across multiple studies with variable designs and healthcare system with different infrastructures and institutional volumes and capacities, underscoring the global variability in emergency CRC presentation and management. A central strength is the explicit focus on **risk stratification and heterogeneity**, illustrating the reason of the constrained of direct comparison of strategies.

Limitations: This review is limited by major heterogeneity in multiple aspects including study design, definitions, physiological instability reporting, and

outcome measures, which precluded meta-analysis and constrained causal inference. Most included studies were observational, with likely confounding by indication and risk selection as higher-risk patients preferentially receiving diversion or Hartmann's procedure. Long-term outcomes such as recurrence, survival, quality of life, and permanent stoma rates, were inconsistently reported, limiting robust conclusions regarding oncologic and functional outcomes.

RECOMMENDATIONS

- For decision making in obstructed and perforated CRC management adopt risk-stratified decision-making by documenting physiological status such as sepsis, hemodynamic stability, contamination severity, and comorbidity burden at presentation.
- Use structured standardized reporting in future studies including definition of emergency presentation, obstruction/perforation subtype, mortality timepoints (30- and/or 90-day), morbidity grading (e.g., Clavien–Dindo), and stoma type (temporary vs permanent).
- Use SEMS as BTS selectively for left-sided obstructing tumors in the absence of perforation/peritonitis and only where appropriate facilities and expertise is available.
- Adopt a pragmatic a perioperative strategy for the emergency management after initial resuscitation including analgesia optimization, fluid strategy, early mobilization when, rather than claiming full ERAS equivalence across unstable cohorts.
- Implement system-level audit and registry protocols to reduce outcome variability and enable relevant benchmarking across institutions.

CONCLUSION

Perforated and obstructed colorectal cancers presenting as emergency are a high-risk surgical condition associated with considerable morbidity and mortality. Evidence generated in this review demonstrates that outcomes are primarily determined by physiological instability, septic burden, and system-level factors, rather than tumor characteristics alone. Perforation with generalized peritonitis carries the greatest early mortality, whereas obstructive disease characterized by a broader

physiological spectrum in which risk is strongly modulated by baseline physiological reserve and the timeliness of resuscitative care. Interpretation of the literature is considerably constrained by heterogeneity in definitions, severity reporting, and outcome measurement, together with confounding by indication in operative strategy selection. Consequently, reported differences between surgical approaches often demonstrate case-mix variation rather than primary treatment superiority. These findings support a physiology-driven approach for interpreting emergency colorectal cancer outcomes, in which the predeterminant risk factors include septic burden, physiological reserve, and system-level factors. Standardization of definitions and outcome documentation is critical to enable reliable comparison across studies and to guide future clinical research.

ABBREVIATIONS:

ASA: American Society of Anesthesiologists

CRC: Colorectal cancer

ERAS: Enhanced Recovery After Surgery

ICU: Intensive care unit

LOS: Length of Stay

MDT: Multidisciplinary team

SEMS: Self-expandable metallic stent

POSSUM: Physiological Operative Severity Score enumeration of Morbidity and Mortality

NSQIP: National Surgical Quality Improvement Program

OS: Overall survival

DFS: disease free survival

CSS: Cancer specific survival

DECLARATION:

Authors contribution:

A.K.E.: Conceptualization, methodology, literature search, writing – original draft.

M.P.K.W.: Writing – review and editing.

I.S.: Supervision, review, and manuscript finalization.

Conflict of interest: The authors declare no conflict of interest.

Funding: No funding was received.

Ethical approval and accordance: Not applicable.

Data availability: Not applicable.

Consent to participate: Not applicable.

Clinical trial number: Not Applicable

REFERENCES

1. Akimoto, N., Ugai, T., Zhong, R., et al. (2021). Rising incidence of early-onset colorectal cancer: A call to action. *Nature Reviews Clinical Oncology*, 18, 230–243. <https://doi.org/10.1038/s41571-020-00445-1>
2. Hofseth, L. J., Hebert, J. R., Chanda, A., et al. (2020). Early-onset colorectal cancer: Initial clues and current views. *Nature Reviews Gastroenterology & Hepatology*, 17, 352–364. <https://doi.org/10.1038/s41575-019-0253-4>
3. Wu, C. W., & Lui, R. N. (2022). Early-onset colorectal cancer: Current insights and future directions. *World Journal of Gastrointestinal Oncology*, 14, 230–241. <https://doi.org/10.4251/wjgo.v14.i1.230>
4. Sung, H., Ferlay, J., Siegel, R. L., et al. (2021). Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians*, 71(3), 209–249. <https://doi.org/10.3322/caac.21660>
5. Kim, M. H., Park, S., Yi, N., et al. (2022). Colorectal cancer mortality trends in the era of cancer survivorship in Korea: 2000–2020. *Annals of Coloproctology*, 38, 343. <https://doi.org/10.3393/ac.2022.00535.0076>
6. Bretthauer, M., Løberg, M., Wieszczyn, P., et al. (2022). Effect of colonoscopy screening on risks of colorectal cancer and related death. *The New England Journal of Medicine*, 387(17), 1547–1556. <https://doi.org/10.1056/NEJMoa2208375>
7. McPhail, S., Johnson, S., Greenberg, D., et al. (2013). Emergency presentation of cancer and short-term mortality. *British Journal of Cancer*, 109, 2027–2034. <https://doi.org/10.1038/bjc.2013.569>
8. Golder, A. M., McMillan, D. C., Horgan, P. G., et al. (2022). Determinants of emergency presentation in patients with colorectal cancer: A systematic review and meta-analysis. *Scientific Reports*, 12, 4366. <https://doi.org/10.1038/s41598-022-08447-8>
9. Bass, G., Fleming, C., Conneely, J., et al. (2009). Emergency first presentation of colorectal cancer predicts significantly poorer outcomes: A review of 356 consecutive Irish patients. *Diseases of the Colon & Rectum*, 52, 678–684. <https://doi.org/10.1007/DCR.0b013e3181a1d8c9>
10. Moura, A. R., Marques, A. D., Dantas, M. S., et al. (2020). Trends in the incidence and mortality of colorectal cancer in a Brazilian city. *BMC Research Notes*, 13, 560. <https://doi.org/10.1186/s13104-020-05411-9>
11. Gunnarsson, H., Ekholm, A., Olsson, L., et al. (2013). Emergency presentation and socioeconomic status in colon cancer. *European Journal of Surgical Oncology*, 39, 831–836. <https://doi.org/10.1016/j.ejso.2013.04.004>
12. Neal, R. D., Tharmanathan, P., France, B., et al. (2015). Is increased time to diagnosis and treatment in symptomatic cancer associated with poorer outcomes? Systematic review. *British Journal of Cancer*, 112(Suppl 1), S92–S107. <https://doi.org/10.1038/bjc.2015.48>
13. Hamilton, W., & Bailey, S. E. (2023). Colorectal cancer in symptomatic patients: How to improve the diagnostic pathway. *Best Practice & Research Clinical Gastroenterology*, 66, 101842. <https://doi.org/10.1016/j.bpg.2023.101842>
14. Pisano, M., Zorcolo, L., Merli, C., et al. (2018). 2017 WSES guidelines on colon and rectal cancer emergencies: Obstruction and perforation. *World Journal of Emergency Surgery*, 13, 36. <https://doi.org/10.1186/s13017-018-0192-3>
15. Yang, K. M., Jeong, M. J., Yoon, K. H., et al. (2022). Oncologic outcome of colon cancer with perforation and obstruction. *BMC Gastroenterology*, 22, 247. <https://doi.org/10.1186/s12876-022-02319-5>
16. Otani, K., Kawai, K., Hata, K., et al. (2019). Colon cancer with perforation. *Surgery Today*, 49(1), 15–20. <https://doi.org/10.1007/s00595-018-1661-8>
17. Norman, J., & Moodley, Y. (2023). Large bowel perforation in patients with colorectal cancer: A South African perspective. *Journal of Cancer & Allied Specialties*, 9, 517. <https://doi.org/10.37029/jcas.v9i1.517>
18. Asano, H., Fukano, H., Takagi, M., & Takayama, T. (2023). Risk factors for recurrence of stage II perforated colorectal cancer: A

- retrospective observational study. *Asian Journal of Surgery*, 46(1), 201–206.
<https://doi.org/10.1016/j.asjsur.2022.03.026>
19. Turri, G., Caliskan, G., Conti, C., et al. (2022). Impact of age and comorbidities on short- and long-term outcomes of patients undergoing surgery for colorectal cancer. *Frontiers in Oncology*, 12, 959650.
<https://doi.org/10.3389/fonc.2022.959650>
20. Krutsri, C., Sumpritpradit, P., Singhatas, P., et al. (2021). Morbidity, mortality, and risk factors of emergency colorectal surgery among older patients in the Acute Care Surgery service: A retrospective study. *Annals of Medicine and Surgery*, 62, 485–489.
<https://doi.org/10.1016/j.amsu.2020.11.001>
21. Mualla, N. M., Hussain, M. R., Akmah, M., Malik, P., Bashir, S., & Lin, J. J. (2021). The impact of postoperative complications on long-term oncological outcomes following curative resection of colorectal cancer (stage I–III): A systematic review and meta-analysis. *Cureus*, 13, e12837.
<https://doi.org/10.7759/cureus.12837>
22. Esswein, K., Ninkovic, M., Gasser, E., et al. (2023). Emergency resection is an independent risk factor for decreased long-term overall survival in colorectal cancer: A matched-pair analysis. *World Journal of Surgical Oncology*, 21, 310.
<https://doi.org/10.1186/s12957-023-03182-8>
23. Arnarson, Ö., Syk, I., Butt, S. T., et al. (2023). Who should operate on patients presenting with emergent colon cancer? A comparison of short- and long-term outcomes depending on surgical subspecialization. *World Journal of Emergency Surgery*, 18, 3.
<https://doi.org/10.1186/s13017-023-00474-y>
24. Sato, K., Fukunaga, Y., Takamatsu, M., et al. (2025). Short- and long-term outcomes of one-stage radical resection and anastomosis without preoperative decompression and diverting stoma between incomplete obstructive and non-obstructive left-sided colorectal cancer: A retrospective observational study. *Journal of the Anus, Rectum and Colon*, 9(1), 41–51.
<https://doi.org/10.23922/jarc.2024-076>
25. van Hooft, J. E., van der Veld, J. V., Arnold, D., et al. (2020). Self-expandable metal stents for obstructing colonic and extracolonic cancer: European Society of Gastrointestinal Endoscopy (ESGE) guideline—Update 2020. *Endoscopy*, 52, 389–407.
<https://doi.org/10.1055/a-1140-3017>
26. Pavlidis, E. T., Galanis, I. N., & Pavlidis, T. E. (2024). Management of obstructed colorectal carcinoma in an emergency setting: An update. *World Journal of Gastrointestinal Oncology*, 16, 598–613.
<https://doi.org/10.4251/wjgo.v16.i3.598>
27. van Hooft, J. E., Bemelman, W. A., Oldenburg, B., et al. (2011). Colonic stenting versus emergency surgery for acute left-sided malignant colonic obstruction: A multicentre randomised trial. *The Lancet Oncology*, 12(4), 344–352. [https://doi.org/10.1016/S1470-2045\(11\)70035-3](https://doi.org/10.1016/S1470-2045(11)70035-3)
28. CReST Collaborative Group. (2022). Colorectal Endoscopic Stenting Trial (CReST) for obstructing left-sided colorectal cancer: Randomized clinical trial. *British Journal of Surgery*, 109(11), 1073–1080.
<https://doi.org/10.1093/bjs/znac141>
29. Binetti, M., Lauro, A., Tonini, V., et al. (2022). Colonic stent as a bridge to surgery for acute left-sided malignant obstruction. *World Journal of Clinical Oncology*, 13, 957–966.
<https://doi.org/10.5306/wjco.v13.i12.957>
30. Ma, W., Zhang, J. C., Luo, K., et al. (2023). Self-expanding metal stents versus decompression tubes as a bridge to surgery for patients with obstruction caused by colorectal cancer: A systematic review and meta-analysis. *World Journal of Emergency Surgery*, 18, 46.
<https://doi.org/10.1186/s13017-023-00515-6>
31. Kim, D. H., & Lee, H. H. (2024). Colon stenting as a bridge to surgery in obstructive colorectal cancer management. *Clinical Endoscopy*, 57, 424–433.
<https://doi.org/10.5946/ce.2023.138>
32. Sloothaak, D. A. M., van den Berg, M. W., Dijkgraaf, M. G. W., et al. (2014). Oncological outcome of malignant colonic obstruction in the Dutch Stent-In 2 trial. *British Journal of Surgery*, 101(13), 1751–1757. <https://doi.org/10.1002/bjs.9645>
33. Arezzo, A., Passera, R., Lo Secco, G., et al. (2017). Colonic stenting as a bridge to surgery versus emergency surgery for malignant colonic obstruction: Results of a multicentre randomized controlled trial (ESCO trial). *Surgical Endoscopy*,

- 31(8), 3297–3305. <https://doi.org/10.1007/s00464-016-5362-3>
34. Alselaime, N. A., Altoub, H. A., Alhassan, M. K., et al. (2022). The role of laparoscopy in emergency colorectal surgery. *Saudi Medical Journal*, 43, 1333–1340. <https://doi.org/10.15537/smj.2022.43.12.20220658>
35. Shida, D., Tagawa, K., Inada, K., et al. (2020). Enhanced recovery after surgery (ERAS) protocols for emergency colorectal surgery: A systematic review and meta-analysis. *International Journal of Colorectal Disease*, 35(2), 225–235. <https://doi.org/10.1007/s00384-019-03476-0>
36. Teixeira, U. F., Goldoni, M. B., Waechter, F. L., Sampaio, J. A., Utiyama, E. M., & Rasslan, S. (2021). Enhanced recovery (ERAS) protocol in emergency colorectal surgery: A prospective cohort study. *International Journal of Colorectal Disease*, 36(5), 1027–1035. <https://doi.org/10.1007/s00384-021-03852-6>
37. Lohsiriwat, V., & Jitmongkarn, R. (2019). Enhanced recovery after surgery in emergency colorectal surgery: Review of literature and current practices. *World Journal of Gastrointestinal Surgery*, 11, 41–52. <https://doi.org/10.4240/wjgs.v11.i2.41>
38. van der Veld, J. V., Amelung, F. J., Borstlap, W. A. A., van Hooft, J. E., Tanis, P. J., & Bemelman, W. A. (2021). Definition of large bowel obstruction by primary colorectal cancer: A systematic review. *Colorectal Disease*, 23(4), 787–804. <https://doi.org/10.1111/codi.15479>
39. Qaderi, S. M., Galjart, B., Verhoef, C., et al. (2021). Disease recurrence after colorectal cancer surgery in the modern era: A population-based study. *International Journal of Colorectal Disease*, 36, 2399–2410. <https://doi.org/10.1007/s00384-021-03914-w>
40. Mikalonis, M., Avlund, T. H., & Løve, U. S. (2024). Danish guidelines for treating acute colonic obstruction caused by colorectal cancer—A review. *Frontiers in Surgery*, 11, 1400814. <https://doi.org/10.3389/fsurg.2024.1400814>
41. Ansaloni, L., Andersson, R. E., Bazzoli, F., et al. (2010). Guidelines in the management of obstructing cancer of the left colon. *World Journal of Emergency Surgery*, 5, 29. <https://doi.org/10.1186/1749-7922-5-29>
42. Yoo, R. N., Cho, H. M., & Kye, B. H. (2021). Management of obstructive colon cancer: Current status, obstacles, and future directions. *World Journal of Gastrointestinal Oncology*, 13, 1850–1862. <https://doi.org/10.4251/wjgo.v13.i12.1850>
43. Grigorean, V. T., Erchid, A., Coman, I. S., & Lițescu, M. (2023). Colorectal cancer—The “parent” of low bowel obstruction. *Medicina*, 59, 875. <https://doi.org/10.3390/medicina59050875>
44. Chen, M., Huang, T., & Wang, C. (2017). Outcome of colon cancer initially presenting as colon perforation and obstruction. *World Journal of Surgical Oncology*, 15, 164. <https://doi.org/10.1186/s12957-017-1228-y>
45. Harji, D. P., Griffiths, B., Stocken, D., et al. (2024). Protocolized care pathways in emergency general surgery: A systematic review and meta-analysis. *British Journal of Surgery*, 111, znae057. <https://doi.org/10.1093/bjs/znae057>
46. Boeding, J. R. E., Elferink, M. A. G., Tanis, P. J., et al. (2023). Surgical treatment and overall survival in patients with right-sided obstructing colon cancer—A nationwide retrospective cohort study. *International Journal of Colorectal Disease*, 38, 248. <https://doi.org/10.1007/s00384-023-04541-3>
47. Delamare Fauvel, A., Bischof, J. J., Reinbolt, R. E., Weihing, V. K., Boyer, E. W., Caterino, J. M., & Wang, H. E. (2023). Diagnosis of cancer in the emergency department: A scoping review. *Cancer Medicine*, 12, 8710–8728. <https://doi.org/10.1002/cam4.5600>
48. Storli, P. E., Dille Amdam, R. G., Skjærseth, G. H., et al. (2023). Cumulative incidence of first recurrence after curative treatment of stage I–III colorectal cancer: Competing risk analyses of temporal and anatomic patterns. *Acta Oncologica*, 62, 1822–1830. <https://doi.org/10.1080/0284186X.2023.2269644>
49. Traiki, T., AlShammari, A., AlRabah, R. N., et al. (2023). Oncological outcomes of elective versus emergency surgery for colon cancer: A tertiary academic center experience. *Saudi Journal of Gastroenterology*, 29, 316–322. https://doi.org/10.4103/sjg.sjg_31_23

50. Acar, N., Acar, T., Kamer, E., et al. (2020). Should we still doubt the success of emergency oncologic colorectal surgery? A retrospective review. *Ulusal Travma ve Acil Cerrahi Dergisi*, 26, 55–62. <https://doi.org/10.14744/tjtes.2019.04043>
51. Elmessiry, M. M., & Mohamed, E. A. (2020). Emergency curative resection of colorectal cancer: Do it with caution. A comparative case series. *Annals of Medicine and Surgery*, 55, 70–76. <https://doi.org/10.1016/j.amsu.2020.04.033>
52. Sandén, G., Svensson, J., Ljuslinder, I., et al. (2023). Defunctioning stoma before neoadjuvant treatment or resection of endoscopically obstructing rectal cancer. *International Journal of Colorectal Disease*, 38, 24. <https://doi.org/10.1007/s00384-023-04318-8>
53. Mauro, A., Scalvini, D., Borgetto, S., Fugazzola, P., Mazza, S., Perretti, I., et al. (2024). Malignant acute colonic obstruction: Multidisciplinary approach for endoscopic management. *Cancers*, 16(4), 821. <https://doi.org/10.3390/cancers16040821>
54. Kim, E. M., Park, J. H., Kim, B. C., et al. (2023). Self-expandable metallic stents as a bridge to surgery in obstructive right- and left-sided colorectal cancer: A multicenter cohort study. *Scientific Reports*, 13, 438. <https://doi.org/10.1038/s41598-023-27767-1>
55. Maertens, V., Stefan, S., Rawlinson, E., et al. (2022). Emergency robotic colorectal surgery during the COVID-19 pandemic: A retrospective case series. *Journal of Laparoendoscopic & Advanced Surgical Techniques. Part A*, 32, 57–60.
56. Prasongdee, R. (2023). Comparison of the short-term outcomes after primary anastomosis between an emergency operation with manual fecal decompression in completely obstructed left-sided colorectal cancer and non-obstructed colorectal cancer in elective bowel preparation: A retrospective single-center study. *Thai Journal of Surgery*, 44, 76–81.
57. Lam, V., Loon, M. M., Alrawe, M., Abougendy, I. S., & Ali, M. (2025). Comparative predictive accuracy of ASA, POSSUM, and NSQIP risk scoring systems in major abdominal surgeries: A systematic review. *Cureus*, 17(6), e85572. <https://doi.org/10.7759/cureus.85572>
58. Yang, B., Lin, J., Deng, Z., et al. (2017). Reviewing the management of obstructive left colon cancer: Assessing the feasibility of one-stage resection and anastomosis after intraoperative colonic irrigation. *Clinical Colorectal Cancer*, 16, e89–e103. <https://doi.org/10.1016/j.clcc.2016.12.001>
59. Kobayashi, H., Asano, M., Ishiguro, M., et al. (2024). Multi-institutional registry of large bowel cancer in Japan conducted by the JSCCR: Cases treated in 2015. *Journal of the Anus, Rectum and Colon*, 8, 265–270. <https://doi.org/10.23922/jarc.2024-065>
60. Wong, N. W., Jabbar, S. A. A., Ngu, J. C. Y., & Teo, N. Z. (2025). Minimally invasive surgery for colorectal cancer emergencies. *World Journal of Clinical Oncology*, 16(8), 107757. <https://doi.org/10.5306/wjco.v16.i8.107757>
61. Sahin, M. K., Yilmaz, B., & Timuroglu, A. (2026). Can the frailty score independently predict postoperative morbidity in patients with colorectal cancer? A prospective observational study. *BMC Geriatrics*, 26, Advance online publication. <https://doi.org/10.1186/s12877-026-07255-7>
62. Wong, T., Pattarapuntakul, T., Netinatsunton, N., Sottisuporn, J., Yaowmanerat, T., Chaochankit, W., et al. (2025). Stent as a bridge to surgery for malignant colonic obstruction: A retrospective study on survival and outcomes. *BMC Gastroenterology*, 25(1), 55. <https://doi.org/10.1186/s12876-025-03654-z>
63. El Hussuna, A., Knudsen, M., Frasson, M., & Poulsen, L. Ø. (2023). Outcomes of emergency surgical interventions in right-sided colonic cancer: Nationwide population-based study based on the Danish Colorectal Cancer Group register. *BJS Open*, 7, zrac153. <https://doi.org/10.1093/bjsopen/zrac153>
64. Zhang, Z., Wu, T., Cai, X., et al. (2024). Feasibility of neoadjuvant chemotherapy for locally advanced, operable colon cancer: Pilot phase of a randomized controlled trial. *eClinicalMedicine*, 63, 102415.
65. Khoo, M. S. Q., Koh, F. H., Sivarajah, S. S., Ho, L. M. L., Aw, D. K. L., Chong, C. X. Z., et al. (2024). Colonic stenting: Is the bridge to surgery worth its cost? A cost-effectiveness analysis at a single Asian institution. *Annals of Coloproctology*, 40(6), 555–563. <https://doi.org/10.3393/ac.2023.00738.0105>