

Macroscopic Normal Appendix in Patients Presented with Features Suggestive of Acute Appendicitis

Salah Eldin Abbas Hamza¹, Ahmed Al-Makki², Amer Abdullah Hamza³

¹Department of Pathology, Soba University Hospital

²Department of Surgery, Khartoum Teaching Hospital

³Department of Surgery, Omdurman Teaching Hospital

Abstract:

Background: Acute appendicitis is one of the most common causes of abdominal pain requiring surgical intervention. However, a subset of patients presents with classical symptoms suggestive of appendicitis, but intraoperative findings reveal a macroscopically normal appendix. This study aimed to evaluate the clinical, radiological, and histopathological characteristics of such cases. **Methods:** This prospective study included 100 patients who presented with clinical features suggestive of acute appendicitis and underwent appendectomy. Only those with macroscopically normal appendices were included. Clinical symptoms, physical signs, laboratory and radiological findings, intraoperative observations, and histopathological results were collected and analyzed. **Results:** Out of 100 patients 65 were females and 35 males. All of them reported right iliac fossa pain. Nausea was present in 80%, fever in 67%, and vomiting in 49%. Tenderness in the right side of the abdomen was noted in 97 patients (69 females, 28 males), while 3 patients (2 females, 1 male) presented with flank pain, anorexia, elevated white blood cell count, and negative β -hCG. Per rectal examination in these 3 patients revealed tenderness. Clinical examination showed RIF tenderness in 56% and rebound tenderness in 41%. Fever was documented in 53%. White blood cell count was elevated in 45% of patients. Ultrasound, performed in female patients only (40 cases), was non-diagnostic in all. Intraoperatively, all appendices were macroscopically normal. Thirteen females were found to have ruptured ovarian follicles. Histopathological examination revealed true appendicitis in 19 patients (13 males and 6 females), including 6 with normal white blood cell counts. **Conclusion:** A macroscopically normal appendix may still harbor histological evidence of inflammation. Clinical judgment should be supported by histopathology, especially in female patients, where gynecological conditions may mimic appendicitis. Improved diagnostic tools and imaging protocols are essential to reduce negative appendectomy rates.

Correspondence to: Salah Eldin Abbas Hamza, salahhamza038@gmail.com

Keywords: Macroscopic Normal Appendix, Acute Appendicitis

Introduction:

Acute appendicitis remains the most common surgical emergency worldwide. Early diagnosis is critical to avoid complications such as perforation and peritonitis. Despite technological advancements in imaging,

diagnosis often relies on clinical judgment, which can be misleading in atypical presentations. A peculiar subgroup of patients presents with classical symptoms, yet intraoperative findings show a macroscopically normal appendix. This raises critical questions:

Should such appendices be removed? Can subclinical inflammation be detected histologically? Are alternative diagnoses, particularly gynecological ones in females, being overlooked?

This study aims to explore the clinical, imaging, intraoperative, and histopathological findings in patients with normal-appearing appendices during surgery for suspected acute appendicitis.

Discussion:

This study highlights the diagnostic dilemma posed by patients presenting with clinical features of acute appendicitis but found intraoperatively to have a macroscopically normal appendix. Despite normal gross findings, histopathological evaluation revealed subclinical appendicitis in 19% of cases, underscoring the importance of routine histological examination even in seemingly normal specimens.

The findings reinforce previous literature indicating that a macroscopically normal appendix can harbor microscopic inflammation or other subtle pathology not evident to the naked eye. This supports the surgical decision to proceed with appendectomy based on clinical suspicion, as withholding surgery based solely on intraoperative appearance could lead to missed diagnoses and potential complications.

Of particular note is the high incidence of gynecological conditions, especially ruptured ovarian follicles, found intraoperatively in female patients. This emphasizes the necessity for comprehensive preoperative gynecologic assessment and better integration of imaging techniques such as transvaginal ultrasound in female patients of reproductive age presenting with right iliac fossa pain.

The study also illustrates the limitations of current diagnostic modalities. Laboratory

markers such as leukocytosis were absent in more than half of histologically confirmed appendicitis cases. Similarly, ultrasound, performed only in female patients, yielded non-diagnostic results, further complicating decision-making.

To reduce the negative appendectomy rate, especially in females, a multidisciplinary approach including enhanced imaging protocols, gynecological consultation, and potentially diagnostic laparoscopy should be considered. Moreover, these findings advocate for the development of clinical scoring systems that integrate gender-specific presentations and non-inflammatory mimics of appendicitis.

The authors would like to express their sincere gratitude to the staff of the Departments of Pathology and Surgery at Soba University Hospital, Khartoum Teaching Hospital, and Omdurman Teaching Hospital for their invaluable support during data collection and patient management. Special thanks go to the laboratory and operating room personnel for their collaboration and dedication throughout this study.

References:

1. Bhangu A, Søreide K, Di Saverio S, Assarsson JH, Drake FT. Acute appendicitis: modern understanding of pathogenesis, diagnosis, and management. *Lancet*. 2015;386(10000):1278–1287.
2. Andersson RE. The natural history and traditional management of appendicitis revisited: spontaneous resolution and predominance of prehospital perforations imply that a correct diagnosis is more important than an early diagnosis. *World J Surg*. 2007;31(1):86–92.
3. Gaitini D. Imaging acute appendicitis: state of the art. *J Clin Imaging Sci*. 2011;1:49.
4. Humes DJ, Simpson J. Acute appendicitis. *BMJ*. 2006;333(7567):530–534.
5. Oyetunji TA, Ong’uti SK, Bolorunduro OB, et al. Pediatric negative appendectomy rate: trend, predictors, and differentials. *J Surg Res*. 2012;173(1):16–20.
6. Kretzschmar K, Tillemans M, Mertens H. A macroscopically normal appendix: is there such a thing? *J Pediatr Surg*. 1990;25(10):1070–1072.